



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\begin{vmatrix} x & y & -z \\ y & -z & x \\ -z & x & y \end{vmatrix}^3 = \begin{vmatrix} X & Y & Z \\ Y & Z & X \\ Z & X & Y \end{vmatrix} \dots (2),$$

where

$$\begin{aligned} X &= x^3 + y^3 - z^3 - 3xyz \\ Y &= 3(x^2y + xz^2 - y^2z) = 0. \\ Z &= 3(xy^2 + yz^2 - x^2z) \end{aligned}$$

Then substituting in (2) and expanding

$$X^3 + Z^3 = (x^3 + y^3 - z^3 + 3xyz)^3$$

which is impossible, since the sum of two integral cubes cannot be an integral cube. (For a proof, see Euler's *Algebra*.) Hence the impossibility of (1) is established.

Also solved by the late *JOSIAH H. DRUMMOND*.

### AVERAGE AND PROBABILITY.

122. Proposed by *F. M. PRIEST*, St. Louis, Mo.

Suppose each of the nine digits to be placed in a wheel, and five of them drawn at random therefrom, and written down in the order drawn. What is the probability the number thus expressed will be greater than 50,000?

Solution by *G. B. M. ZERR*, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa., and *J. F. LAWRENCE*, A. B., Breckenridge, Mo.

If 5, 6, 7, 8, or 9 be drawn first, the number will be greater than 50,000.


The chance of drawing 5 is  $\frac{1}{9}$ ; of drawing 6,  $\frac{1}{9}$ ; of drawing 7,  $\frac{1}{9}$ ; of drawing 8,  $\frac{1}{9}$ ; of drawing 9,  $\frac{1}{9}$ . The chance of drawing 5, 6, 7, 8, or 9 is, therefore,  $\frac{5}{9}$ . Therefore the chance that the number is greater than 50,000 is  $\frac{5}{9}$ .

123. Proposed by *L. C. WALKER*, A. M., Graduate Student, Leland Stanford Jr. University, Cal.

Three points are taken at random within a square. What is the probability that the angle formed by joining them is acute?

Solution by *G. B. M. ZERR*, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Let  $ABCD$  be the square;  $P, Q, R$  the three random points. Through  $P, Q$  draw  $GH$ , and through  $Q$  draw  $KL$  perpendicular to  $GH$ . When  $P$  is between  $G$  and  $Q$  the angle  $PQR$  will be obtuse if  $R$  lies on the opposite side of  $LK$  from  $P$ . Draw  $AF$  perpendicular to  $KL$ ;  $AE$  perpendicular to  $GH$ ;  $ST$  parallel to  $GH$ ;  $MN$  and  $TU$  parallel to  $KL$ ;  $DM$  parallel to  $GH$ .



Let  $AB=a$ ,  $PQ=x$ ,  $AF=y$ ,  $AE=z$ ,  $\angle KAF=\theta$ . Then  $AK=y\sec\theta$ ,  $BK=$

$a - y \sec \theta$ , when  $L$  is on  $BC$ ,  $BL = (a - y \sec \theta) \cot \theta$ , when  $L$  is on  $CD$ , and  $CL = a(1 - \tan \theta) - y \sec \theta$ . Area  $L\dot{B}K = \frac{1}{2}(a - y \sec \theta)^2 \cot \theta = \frac{1}{2}u$ . Area  $L\dot{O}BK = \frac{1}{2}a(2a - 2y \sec \theta - a \tan \theta) = \frac{1}{2}u_1$ . An element of area at  $P$  is  $x dx d\theta$ ; at  $Q$ ,  $dy dz$ . The limits of  $\theta$  for  $\frac{1}{2}u$  are  $\frac{1}{4}\pi$  and  $\frac{1}{2}\pi$ , and 0 and  $\frac{1}{4}\pi$ ; for  $\frac{1}{2}u_1$ , 0 and  $\frac{1}{4}\pi$ ; of  $y$  for  $\theta = \frac{1}{4}\pi$  to  $\theta = \frac{1}{2}\pi$ ,  $y = 0$  to  $y = a \cos \theta$ ; for  $\theta = 0$  to  $\theta = \frac{1}{4}\pi$ ,  $y = AN = a(\sec \theta - \sin \theta) = y_2$  to  $y = a \cos \theta$ , and  $y = AU = a(1 - \tan \theta) \sec \theta = y_3$  to  $y = y_2$ , and  $y = a(\cos \theta - \sin \theta) = y_1$  to  $y = y_3$ , for  $\frac{1}{2}u_1$ ,  $y = 0$  to  $y = y_1$ ; of  $z$ , for  $y = 0$  to  $a \cos \theta$ ,  $z = y \tan \theta$  to  $z = a \cos \theta$  and  $z = a \cos \theta$  to  $z = a \operatorname{cosec} \theta - y \cot \theta = z_1$ ; for  $y = y_2$  to  $y = a \cos \theta$ ,  $z = y \tan \theta$  to  $z = z_1$ , for  $y = y_3$  to  $y = y_2$ ,  $z = y \tan \theta$  to  $z = a \cos \theta$  and  $z = a \cos \theta$  to  $z = z_1$ , for  $y = y_1$  to  $y = y_3$ ,  $z = y \tan \theta$  to  $z = a \cos \theta$  and  $z = a \cos \theta$  to  $z = a \sec \theta$ , for  $\frac{1}{2}u_1$  the limits of  $z$  are  $y \tan \theta$  to  $a \cos \theta$  and  $a \cos \theta$  to  $a \sec \theta$ ; of  $x$ , 0 and  $y + z \tan \theta = x_1$ , and 0 and  $y + a \operatorname{cosec} \theta - z \cot \theta = x_2$ . The limits of  $x$  must be doubled for the case when  $P$  is on the opposite side of  $LK$ . The whole number of ways the three points can be taken is  $a^6$ . Let  $p'$  be the chance that  $PQR$  is obtuse.

$$\begin{aligned}
 p' = & \frac{1}{a^6} \left[ \int_{\frac{1}{4}\pi}^{\frac{1}{2}\pi} \int_0^{a \cos \theta} u d\theta du + \int_0^{\frac{1}{4}\pi} \int_{y_3}^{y_2} u d\theta du \right] \left[ \int_{y \tan \theta}^{x_1} \int_0^{x_2} dz dx + \int_{a \cos \theta}^{z_1} \int_0^{x_2} dz dx \right] \\
 & + \frac{1}{a^6} \left[ \int_0^{\frac{1}{4}\pi} \int_{y_1}^{y_3} u d\theta du + \int_0^{\frac{1}{4}\pi} \int_0^{y_1} u_1 d\theta du \right] \left[ \int_{y \tan \theta}^{x_1} \int_0^{x_2} dz dx + \int_{a \cos \theta}^{a \sec \theta} \int_0^{x_2} dz dx \right] \\
 & + \frac{1}{a^6} \int_0^{\frac{1}{4}\pi} \int_{y_2}^{a \cos \theta} \int_{y \tan \theta}^{x_1} u d\theta dy dz dx = \frac{1}{6a^6} \int_{\frac{1}{4}\pi}^{\frac{1}{2}\pi} \int_0^{a \cos \theta} [(y + a \sin \theta)^3 \sec \theta \operatorname{cosec} \theta - \\
 & \quad (y + a \sin \theta - a \cos \theta)^3 \operatorname{cosec}^5 \theta \sec \theta - y^3 \sec^5 \theta \operatorname{cosec} \theta] u d\theta dy \\
 & + \frac{1}{6a^6} \int_0^{\frac{1}{4}\pi} \left[ \int_{y_3}^{y_2} [(y + a \sin \theta)^3 \sec \theta \operatorname{cosec} \theta - y^3 \sec^5 \theta \operatorname{cosec} \theta - (y + a \sin \theta - a \cos \theta)^3 \right. \\
 & \quad \left. \operatorname{cosec}^5 \theta \sec \theta] u dy \right. \\
 & + \int_{y_1}^{y_3} [(y + a \sin \theta)^3 \sec \theta \operatorname{cosec} \theta - y^3 \sec^5 \theta \operatorname{cosec} \theta - y^3 \sin \theta \sec \theta] u dy \\
 & + \int_0^{y_1} [(y + a \sin \theta)^3 \sec \theta \operatorname{cosec} \theta - y^3 \sec^5 \theta \operatorname{cosec} \theta - y^3 \sin \theta \sec \theta] u_1 dy \\
 & \left. + \int_{y_2}^{a \cos \theta} (a^3 \sec^2 \theta \operatorname{cosec} \theta - y^3 \sec^5 \theta \operatorname{cosec} \theta) u dy \right] d\theta \\
 = & \frac{1}{3 \frac{1}{6} 6} \int_{\frac{1}{4}\pi}^{\frac{1}{2}\pi} (14 \sin \theta \cos \theta + 14 \cos^2 \theta + 6 \cot \theta + \cot^2 \theta - \operatorname{cosec}^2 \theta - 20 \cot \theta \operatorname{cosec}^2 \theta \\
 & + 45 \cot^2 \theta \operatorname{cosec}^2 \theta - 36 \cot^3 \theta \operatorname{cosec}^2 \theta + 10 \cot^4 \theta \operatorname{cosec}^2 \theta) d\theta
 \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{360} \int_0^{4\pi} (4\sin^2\theta + 2\cos^2\theta + 42\sin\theta\cos\theta - 12\sin\theta\cos^3\theta - 20\sin^3\theta\cos\theta + \sec^8\theta\sec^2\theta \\
& \quad + 6\sec^7\theta\csc\theta - 60\sec^2\theta\csc\theta - 6\sec\theta\csc\theta - 36\tan\theta + 10\tan^2\theta \\
& \quad + 20\tan^3\theta - \cot^2\theta + 20\csc^2\theta + 123\sec^2\theta - 50\tan\theta\sec^2\theta + 5\tan^2\theta\sec^2\theta \\
& \quad + 14\tan^3\theta\sec^2\theta - 103\tan^4\theta\sec^2\theta + 186\tan^5\theta\sec^2\theta - 236\tan^6\theta\sec^2\theta \\
& \quad + 192\tan^7\theta\sec^2\theta - 60\tan^8\theta\sec^2\theta + 36\tan^9\theta\sec^2\theta + 84\tan^{10}\theta\sec^2\theta \\
& \quad + 45\tan^{12}\theta\sec^2\theta) d\theta = \frac{1}{400} \frac{9}{4} \frac{9}{4} \frac{7}{4} \frac{7}{4} - \frac{1}{144} \log 2.
\end{aligned}$$

$$3p' = \frac{3}{4} \frac{5}{0} \frac{7}{0} \frac{8}{4} \frac{1}{0} - \frac{1}{8} \log 2, \quad p = 1 - 3p' = \frac{1}{4} \frac{1}{8} \log 2 + \frac{4}{0} \frac{0}{0} \frac{6}{4} \frac{1}{0} \frac{9}{0}. \quad \therefore p = .260292.$$

124. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

Find the average area of a spherical polygon of  $n=6$  sides.

No solution of this problem has been received.

### MISCELLANEOUS.

118. Proposed by O. W. ANTHONY, New York, N. Y.

If  $f$  is determined by the equation  $f(\mu\nu) = f(\mu)f^{-1}(\nu) + f(\nu)f^{-1}(\mu)$ , where  $f^{-1}$  is the inverse of  $f$ , show that  $f[(2)^\mu] = \frac{k^{\mu+1}}{2^{\mu+1}}$ , where  $k$  is a constant.

Solution by G. B. M. ZERR, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

$$f(\mu\nu) = ff^{-1}(\mu\nu) + ff^{-1}(\mu\nu), \text{ but } ff^{-1} = 1. \quad \therefore f(\mu\nu) = 2(\mu\nu). \quad \therefore f = 2.$$

$$\therefore (2)^\mu = (f)^\mu \text{ or } f[(2)^\mu] = (f)^\mu + 1 = (2)^{\mu+1}.$$

$$\therefore f[(2)^\mu] = (\frac{1}{2}k)^{\mu+1}, \text{ where } k=4.$$

119. Proposed by L. C. WALKER, A. M., Graduate Student, Leland Stanford Jr. University, Cal.

Show how to determine the illumination at any point of the surface of the water at the bottom of a deep well, due to the light from the sky.

A solution of this problem appeared in the November number. The problem was incorrectly numbered. Ed.

120. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

$$\text{Prove } \Sigma \cos^4 x - 2\Pi \cos^2 x + 2\Pi \sin^2 x = 1 - \sin(\Sigma) \sin \Pi(y+z-x).$$

Solution by G. B. M. ZERR, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

$$\sin(\Sigma) \sin \Pi(y+z-x)$$

$$= \sin(x+y+z) \sin(x+y-z) \sin(z-y+x) \sin(z+y-x)$$